

Day 000 Clinical

Record ID

(REDCap auto generated)



Pharyngeal Electrical stimulation for Acute Stroke dysphagia Trial (PhEAST)

UK ISRCTN 98886991
UK IRAS306761
UK CPMS 50913
WHO UTN U1111-1273-9942

Baseline form (Clinical) v1.7

(Dysphagia/swallow/feeding)

- ▶ Please check consent form obtained.
- ▶ Please check Eligibility form completed.
- ▶ Please check Baseline form completed.

Section A: Participant details

A1. Centre name:

(Centre)

A2. Participant ID :

(Participant ID)

A3. Participant initials (e.g. ABC or A-C) :

(3 uppercase letters, or 2 separated by a hyphen (-))

Section B: Dysphagia Severity Rating Scale (DSRS) at time of enrolment.

DSRS supervision score 3 is always chosen when a patient is on limited or consistent oral trials and still requires NG/PEG/ RIG tube.

B1. DSRS, fluids

- 0: Thin fluids / IDDSI level 0
 - 1: Slightly or mildly thick / IDDSI level 1 or 2
 - 2: Moderately thick / IDDSI level 3
 - 3: Extremely thick / IDDSI level 4
 - 4: No oral fluids
- (Choose one answer)

B2. DSRS, diet

- 0: Regular diet / IDDSI level 7
 1: Easy to chew diet / IDDSI level 8
 2: Soft & bite sized diet / IDDSI level 6
 3: Pureed or minced/moist diet / IDDSI level 4 or 5
 4: No oral feeding
 (Choose one answer)

B3. DSRS, supervision

- 0: Eating independently
 1: Eating with supervision
 2: Feeding by third party (untrained)
 3: Therapeutic feeding (SALT/trained staff)
 4: No oral feeding
 (Choose one answer)

B4. DSRS, total

 (Calculated DSRS)

Section C: Feeding information (NGT/PEG/RIG, FOIS & FSS & IDD FDS & PRESS)

C1. Feeding tube type

- Nasogastric tube (NGT) feeding
 Percutaneous endoscopic gastrostomy (PEG or RIG) tube feeding

C2. Functional oral intake scale (FOIS)

- 1: Nothing by mouth (B4. DSRS=12)
 2: Tube dependent with minimal attempts of food or liquid. (B4. DSRS=11 & B3. DSRS supervision: Therapeutic feeding)
 3: Tube dependent with consistent oral intake of food or liquid
 4: Total oral diet of a single consistency
 5: Total oral diet with multiple consistencies, but requiring special preparation or compensations
 6: Total oral diet with multiple consistencies without special preparation, but with specific food limitations
 7: Total oral diet with no restrictions
 0: Died
 (FOIS Calculated based on DSRS score: [fois_calc_000])

Stroke date [date_stroke] (day [age_stroke_000])

Eligibility: FOIS score of 1 or 2 or FOIS 3

FOIS calculated based on DSRS score: [fois_calc_000]

C2b. FOIS total score

 (Calculated)

NB: FOIS = 0 for Death

C3. Feeding status score (FSS)

- 1: Oral diet/fluids, normal consistency
 2: Oral diet/fluids, soft consistency
 3: Non-oral diet/fluids, nasogastric tube (NGT) feeding
 4: Non-oral diet/fluids, percutaneous endoscopic gastrostomy (PEG or RIG) tube feeding
 5: Non-oral diet/fluids, intravenous/subcutaneous fluids only
 6: No feeding/fluids
 0: Died
 (FSS calculated based on NGT & DSRS score: [fss_calc_000])

FSS calculated based on NGT & DSRS score: [fss_calc_000]

C3b. FSS total score _____

NB: FSS = 0 for Death

△ FOIS/ FSS/ Feeding tube details C1-C3 should all be specified.

IDDSI Functional Diet Scale score



Fig 1 The IDDSI framework.

C4a. Food prescription
(7, 6, 5 are transitional foods)

- 7: REGULAR
- 6: SOFT & BITE-SIZED
- 5: MINCED & MOIST
- 4: PUREED
- 3: LIQUIDISED
- N/A: NO FOOD (N/A)

C4b. Drink prescription

- 4: EXTREMELY THICK
- 3: MODERATELY THICK
- 2: MILDLY THICK
- 1: SLIGHTLY THICK
- 0: THIN
- N/A: NO DRINKS (N/A)

IDDSI Functional Diet Scale score

How is IDDS Scored:

(Calculated)

- If there is an NG Tube or PEG in, with no trials of food and drink (i.e. a FOIS score of 1) score 0
- If there is an NG Tube or PEG in, with trials of food and drink with minimal spoons < 15 (i.e. a FOIS score of 2) score 0.25
- If there is an NG Tube or PEG in, with trials of food and drink with some spoons ≥ 15 (i.e. a FOIS score of 3) score 0.50

- Otherwise, score from the range of 0 to 8 according to the extended IDDSI framework.

Free water protocol DSRS, FOIS, FSS and IDDSI: Based on what the patient is actually doing (i.e. unlimited thin fluids).
Please only complete if the patient is on free water protocol

C5a. Is the participant on a free water protocol? If yes, please complete the DSRS, FOIS, FSS and IDDS below based on what the patient is actually doing (i.e. unlimited thin fluids).

- Yes
 No

Free water protocol DSRS: Based on what the patient is actually doing (i.e. unlimited thin fluids).

C5b. Free water protocol DSRS, fluids Based on what the patient is actually doing (i.e. unlimited thin fluids).

- 0: Thin fluids / IDDSI level 0
 1: Slightly or mildly thick / IDDSI level 1 or 2
 2: Moderately thick / IDDSI level 3
 3: Extremely thick / IDDSI level 4
 4: No oral fluids
 (Choose one answer - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

C5c. Free water protocol DSRS, diet Based on what the patient is actually doing

- 0: Regular diet / IDDSI level 7
 1: Easy to chew diet / IDDSI level 7
 2: Soft & bite sized diet / IDDSI level 6
 3: Pureed or minced/moist diet / IDDSI level 4 or 5
 4: No oral feeding
 (Choose one answer - Based on what the patient is actually doing.)

C5d. Free water protocol DSRS, supervision Based on what the patient is actually doing

- 0: Eating independently
 1: Eating with supervision
 2: Feeding by third party (untrained)
 3: Therapeutic feeding (SALT/trained staff)
 4: No oral feeding
 (Choose one answer - Based on what the patient is actually doing.)

C5e. Free water protocol DSRS, total
Based on what the patient is actually doing (i.e. unlimited thin fluids).

(Calculated DSRS - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

Free water protocol FOIS, FSS: Based on what the patient is actually doing (i.e. unlimited thin fluids).

C5f. Free water protocol Functional oral intake scale (FOIS)

Based on what the patient is actually doing (i.e. unlimited thin fluids).

- 1: Nothing by mouth (B4. DSRS=12)
 - 2: Tube dependent with minimal attempts of food or liquid. (B4. DSRS=11 & B3. DSRS supervision: Therapeutic feeding)
 - 3: Tube dependent with consistent oral intake of food or liquid
 - 4: Total oral diet of a single consistency
 - 5: Total oral diet with multiple consistencies, but requiring special preparation or compensations
 - 6: Total oral diet with multiple consistencies without special preparation, but with specific food limitations
 - 7: Total oral diet with no restrictions
 - 0: Died
- (Choose one answer - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

C5g. Free water protocol Feeding status score (FSS)

Based on what the patient is actually doing (i.e. unlimited thin fluids).

- 1: Oral diet/fluids, normal consistency
 - 2: Oral diet/fluids, soft consistency
 - 3: Non-oral diet/fluids, nasogastric tube (NGT) feeding
 - 4: Non-oral diet/fluids, percutaneous endoscopic gastrostomy (PEG or RIG) tube feeding
 - 5: Non-oral diet/fluids, intravenous/subcutaneous fluids only
 - 6: No feeding/fluids
 - 0: Died
- (Choose one answer - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

Free water protocol IDDSI: Based on what the patient is actually doing (i.e. unlimited thin fluids).

C5h. Free water protocol Food prescription (7, 6, 5 are transitional foods)

Based on what the patient is actually doing (i.e. unlimited thin fluids).

- 7: REGULAR
 - 6: SOFT & BITE-SIZED
 - 5: MINCED & MOIST
 - 4: PUREED
 - 3: LIQUIDISED
 - N/A: NO FOOD (N/A)
- (Choose one answer - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

C5i. Free water protocol Drink prescription

Based on what the patient is actually doing (i.e. thin fluids).

- 4: EXTREMELY THICK
 - 3: MODERATELY THICK
 - 2: MILDLY THICK
 - 1: SLIGHTLY THICK
 - 0: THIN
 - N/A: NO DRINKS (N/A)
- (Choose one answer - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

Free water protocol IDDSI Functional Diet Scale score

Based on what the patient is actually doing (i.e. unlimited thin fluids). How is IDDS Scored:

- If there is an NG Tube or PEG in, with no trials of food and drink (i.e. a FOIS score of 1) score 0
- If there is an NG Tube or PEG in, with trials of food and drink with minimal spoons < 15 (i.e. a FOIS score of 2) score 0.25
- If there is an NG Tube or PEG in, with trials of food and drink with some spoons >=15 (i.e. a FOIS score of 3) score 0.50

- Otherwise, score from the range of 0 to 8 according to the extended IDDSI framework.

(Calculated - Based on what the patient is actually doing (i.e. unlimited thin fluids).)

Predictive Swallowing Score (PRESS)

C6a. Calculated PRESS score. Prognostic Model of Swallowing Recovery and Enteral Tube Feeding After Ischemic Stroke (Galovic et al. JAMA Neuro 2019; 11 Feb)

(Calculated)

C6b. Risk estimate % of impaired oral intake after 7 days

Prediction estimates of Swallowing Recovery According to PRESS value [press_calc_000]

(Risk of Impaired Oral Intake After 7 d. %)

C6c. Risk estimate % of impaired oral intake after 30 days

Prediction estimates of Swallowing Recovery According to PRESS value [press_calc_000]

(Risk of Impaired Oral Intake After 30 d. %)

C6d. Risk estimate % of NO return to prestroke diet after 7 days

Prediction estimates of Swallowing Recovery According to PRESS value [press_calc_000]

(Risk of No Return to Prestroke Diet After 7 d. %)

C6e. Risk estimate % of NO return to prestroke diet after 30 days

Prediction estimates of Swallowing Recovery According to PRESS value [press_calc_000]

(Risk of No Return to Prestroke Diet After 30 d. %)

Section D: EAT-10 swallowing screening tool at time of enrolment.**0= No problem - 5= Not eating**

D1. My swallowing problem has caused me to lose weight?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D2. My swallowing problem interferes with my ability to go out for meals?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D3. Swallowing liquids takes extra effort?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D4. Swallowing solids takes extra effort?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D5. Swallowing pills takes extra effort?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (ask the nurse)

D6. Swallowing is painful?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D7. The pleasure of eating is affected by my swallowing?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D8. When I swallow food sticks in my throat?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D9. I cough when I eat?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (ask the nurse)

D10. Swallowing is stressful?

- No problem
 - Mild
 - Moderate
 - Moderate-severe
 - Severe problem
 - Not eating, i.e. tube fed
- (Choose one answer)

D11a. EAT-10 total score

(Calculated EAT-10 (0-50))

D11b. Average EAT-10 score

(Calculated average EAT-10 (0-50))

△ EAT-10 details D1-D10 should all be specified.

Section E: Modified Rankin Scale (mRS) at time of enrolment.

E1. Modified Rankin Scale (mRS), now

Premorbid / pre-stroke: [mrs_premorbid]

- 0: No symptoms at all
 - 1: No significant disability despite symptoms; able to carry out all usual duties and activities
 - 2: Slight disability; unable to carry out all previous activities, but able to look after own affairs without assistance
 - 3: Moderate disability; requiring some help, but able to walk without assistance
 - 4: Moderately severe disability; unable to walk and attend to bodily needs without assistance
 - 5: Severe disability; bedridden, incontinent and requiring constant nursing care and attention
 - 6: Deceased
 - 7: Withdrawn
(Choose one answer)
-

E2. mRS total score

 (Calculated mRS (0-6))

Section F: Barthel index

F1. Is the participant incontinent of urine?

- Yes, incontinent/has a catheter fitted
 - Occasional accident (maximum once per 24 hours)
 - No, continent
-

F2. How does the participant move from bed to the chair?

- Not at all
 - With a lot of help from one or two people
 - With a little help from one person
 - On their own
-

F3. How does the participant get about?

- Not at all
 - Propelling themselves independently in a wheelchair
 - Walking with the help and supervision of one person
 - Walking with no help (even if they use a stick/frame)
-

F4. Is the participant incontinent of their bowels?

- Yes, incontinent
 - Occasional accident (once per week)
 - No, continent
-

F5. Does the participant wash their own face, brush their teeth and hair (for men, shave)?

- With help
 - Without help
-

F6. How does the participant use the toilet (or commode)?

- With a lot of help
 - With a little help
 - On their own
-

F7. Does the participant feed themselves?

- With major help
 - With some help e.g. cutting
 - Without any help
-

F8. Does the participant need any help with dressing?

- Yes, they need help with almost everything
- Yes, they are able to do about half unaided
- No, they can do everything

F9. How does the participant get up and down the stairs?

- Not at all
 With help (either supervision or assistance)
 Without any help

F10. Does the participant need help with bathing or showering?

- Yes
 No

F11. Barthel index

_____ (Calculated)

△ Barthel index details F1-F11 should be specified.

Section G: National Institutes of Health Stroke Scale (NIHSS)

NIHSS 1a. Level of Consciousness (LOC)

Score 0-1-2: Must be alert (score 0), arouse to minor stimulation (score 1) or require repeated stimulation (score 2) to be eligible.

Score 2-3: Patients with only movements to pain (also score 2) or postures/unresponsive (score 3) are ineligible.

Score 3: The investigator must choose a response if a full evaluation is prevented by such obstacles as an endotracheal tube, language barrier, orotracheal trauma/bandages. A 3 is scored only if the patient makes no movement (other than reflexive posturing) in response to noxious stimulation.

G1a. NIHSS 1a. Level of Consciousness. (from Eligibility form)

- 0 = Alert, keenly responsive
 1 = Arouses to minor stimulation
 2 = Requires repeated stimulation to arouse
 2 = Movements to pain
 3 = Postures or unresponsive
 (Specified on the Eligibility form)

NIHSS 1b. LOC Questions

What is the month and what is your age? The answer must be correct - there is no partial credit for being close. Score 2: Aphasic and stuporous patients who do not comprehend the questions will score 2. Score 1: Patients unable to speak because of endotracheal intubation, orotracheal trauma, severe dysarthria from any cause, language barrier, or any other problem not secondary to aphasia are given a 1. It is important that only the initial answer be graded and that the examiner not "help" the patient with verbal or non-verbal cues.

G1b. NIHSS 1b. What is the month and what is your age?

- 0 = Answers both questions correctly
 1 = Answers one question correctly
 2 = Answers neither question correctly
 (Choose one answer)

NIHSS 1c. LOC Commands

Open and close your eyes. Grip and release your hand. The patient is asked to open and close the eyes and then to grip and release the non-paretic hand. Substitute another one step command if the hands cannot be used. Credit is given if an unequivocal attempt is made but not completed due to weakness. If the patient does not respond to command, the task should be demonstrated to him or her (pantomime), and the result scored (i.e., follows none, one or two commands). Patients with trauma, amputation, or other physical impediments should be given suitable one-step commands. Only the first attempt is scored.

G1c. NIHSS 1c. Open and close your eyes. Grip and release your hand.

- 0 = Performs both tasks correctly
 1 = Performs one task correctly
 2 = Performs neither task correctly
 (Choose one answer)

NIHSS 2. Best gaze.

Only horizontal eye movements will be tested. Voluntary or reflexive (oculocephalic) eye movements will be scored, but caloric testing is not done. If the patient has a conjugate deviation of the eyes that can be overcome by voluntary or reflexive activity, the score will be 1. If a patient has an isolated peripheral nerve paresis (CN III, IV or VI), score a 1. Gaze is testable in all aphasic patients. Patients with ocular trauma, bandages, pre-existing blindness, or other disorder of visual acuity or fields should be tested with reflexive movements, and a choice made by the investigator. Establishing eye contact and then moving about the patient from side to side will occasionally clarify the presence of a partial gaze palsy.

G2. NIHSS 2. Best gaze.

- 0 = Normal
 1 = Partial gaze palsy; gaze is abnormal in one or both eyes, but forced deviation or total gaze paresis is not present
 2 = Forced deviation, or total gaze paresis not overcome by the oculocephalic maneuver
 (Choose one answer)

NIHSS 3. Visual field loss (upper and lower quadrants).

Visual fields (upper and lower quadrants) are tested by confrontation, using finger counting or visual threat, as appropriate. Patients may be encouraged, but if they look at the side of the moving fingers appropriately, this can be scored as normal. If there is unilateral blindness or enucleation, visual fields in the remaining eye are scored. Score 1 only if a clear-cut asymmetry, including quadrantanopia, is found. If patient is blind from any cause, score 3. Double simultaneous stimulation is performed at this point. If there is extinction, patient receives a 1, and the results are used to respond to item 11.

G3. NIHSS 3. Visual field loss (upper and lower quadrants).

- 0 = No visual loss.
 1 = Partial hemianopia.
 2 = Complete hemianopia.
 3 = Bilateral hemianopia (blind including cortical blindness).
 (Choose one answer)

NIHSS 4. Facial palsy.

Ask - or use pantomime to encourage - the patient to show teeth or raise eyebrows and close eyes. Score symmetry of grimace in response to noxious stimuli in the poorly responsive or non-comprehending patient. If facial trauma/bandages, orotracheal tube, tape or other physical barriers obscure the face, these should be removed to the extent possible.

G4. NIHSS 4. Facial palsy.

- 0 = Normal symmetrical movements.
 - 1 = Minor paralysis (flattened nasolabial fold, asymmetry on smiling).
 - 2 = Partial paralysis (total or near-total paralysis of lower face).
 - 3 = Complete paralysis of one or both sides (absence of facial movement in the upper and lower face).
- (Choose one answer)

NIHSS 5. Motor Arm.

The limb is placed in the appropriate position: extend the arms (palms down) 90 degrees (if sitting) or 45 degrees (if supine). Drift is scored if the arm falls before 10 seconds. The aphasic patient is encouraged using urgency in the voice and pantomime, but not noxious stimulation. Each limb is tested in turn, beginning with the non-paretic arm. Only in the case of amputation or joint fusion at the shoulder, the examiner should record the score as untestable (UN), and clearly write the explanation for this choice.

G5a. NIHSS 5a. Motor weakness, left arm

- 0 = No drift; limb holds 90 (or 45) degrees for full 10 seconds.
 - 1 = Drift; limb holds 90 (or 45) degrees, but drifts down before full 10 seconds; does not hit bed or other support.
 - 2 = Some effort against gravity; limb cannot get to or maintain (if cued) 90 (or 45) degrees, drifts down to bed, but has some effort against gravity.
 - 3 = No effort against gravity; limb falls.
 - 4 = No movement.
 - UN = Amputation or joint fusion
- (Choose one answer)

G5b. NIHSS 5b. Motor weakness, right arm

- 0 = No drift; limb holds 90 (or 45) degrees for full 10 seconds.
 - 1 = Drift; limb holds 90 (or 45) degrees, but drifts down before full 10 seconds; does not hit bed or other support.
 - 2 = Some effort against gravity; limb cannot get to or maintain (if cued) 90 (or 45) degrees, drifts down to bed, but has some effort against gravity.
 - 3 = No effort against gravity; limb falls.
 - 4 = No movement.
 - UN = Amputation or joint fusion
- (Choose one answer)

NIHSS 6. Motor Leg.

The limb is placed in the appropriate position: hold the leg at 30 degrees (always tested supine).

Drift is scored if the leg falls before 5 seconds. The aphasic patient is encouraged using urgency in the voice and pantomime, but not noxious stimulation.

Each limb is tested in turn, beginning with the non-paretic leg. Only in the case of amputation or joint fusion at the hip, the examiner should record the score as untestable (UN), and clearly write the explanation for this choice.

G6a. NIHSS 6a. Motor weakness, left leg

- 0 = No drift; leg holds 30-degree position for full 5 seconds.
 - 1 = Drift; leg falls by the end of the 5-second period but does not hit bed.
 - 2 = Some effort against gravity; leg falls to bed by 5 seconds, but has some effort against gravity.
 - 3 = No effort against gravity; leg falls to bed immediately.
 - 4 = No movement.
 - UN = Amputation or joint fusion.
- (Choose one answer)

G6b. NIHSS 6b. Motor weakness, right leg

- 0 = No drift; leg holds 30-degree position for full 5 seconds.
 - 1 = Drift; leg falls by the end of the 5-second period but does not hit bed.
 - 2 = Some effort against gravity; leg falls to bed by 5 seconds, but has some effort against gravity.
 - 3 = No effort against gravity; leg falls to bed immediately.
 - 4 = No movement.
 - UN = Amputation or joint fusion.
- (Choose one answer)

NIHSS 7. Limb Ataxia.

This item is aimed at finding evidence of a unilateral cerebellar lesion. Test with eyes open. In case of visual defect, ensure testing is done in intact visual field. The finger-nose-finger and heel-shin tests are performed on both sides, and ataxia is scored only if present out of proportion to weakness. Ataxia is absent in the patient who cannot understand or is paralyzed. Only in the case of amputation or joint fusion, the examiner should record the score as untestable (UN), and clearly write the explanation for this choice. In case of blindness, test by having the patient touch nose from extended arm position.

G7. NIHSS 7. Limb Ataxia

- 0 = Absent.
 - 1 = Present in one limb.
 - 2 = Present in two limbs.
 - UN = Amputation or joint fusion.
- (Choose one answer)

NIHSS 8. Sensory loss.

Sensation or grimace to pinprick when tested, or withdrawal from noxious stimulus in the obtunded or aphasic patient. Only sensory loss attributed to stroke is scored as abnormal and the examiner should test as many body areas (arms [not hands], legs, trunk, face) as needed to accurately check for hemisensory loss. A score of 2, "severe or total sensory loss," should only be given when a severe or total loss of sensation can be clearly demonstrated. Stuporous and aphasic patients will, therefore, probably score 1 or 0. The patient with brainstem stroke who has bilateral loss of sensation is scored 2. If the patient does not respond and is quadriplegic, score 2. Patients in a coma (item 1a=3) are automatically given a 2 on this item.

G8. NIHSS 8. Sensory loss

- 0 = Normal; no sensory loss.
 - 1 = Mild-to-moderate sensory loss; patient feels pinprick is less sharp or is dull on the affected side; or there is a loss of superficial pain with pinprick, but patient is aware of being touched.
 - 2 = Severe to total sensory loss; patient is not aware of being touched in the face, arm, and leg.
- (Choose one answer)

NIHSS 9. Best language.

A great deal of information about comprehension will be obtained during the preceding sections of the examination. For this scale item, the patient is asked to describe what is happening in the attached picture, to name the items on the attached naming sheet and to read from the attached list of sentences. Comprehension is judged from responses here, as well as to all of the commands in the preceding general neurological exam. If visual loss interferes with the tests, ask the patient to identify objects placed in the hand, repeat, and produce speech. The intubated patient should be asked to write. The patient in a coma (item 1a=3) will automatically score 3 on this item. The examiner must choose a score for the patient with stupor or limited cooperation, but a score of 3 should be used only if the patient is mute and follows no one-step commands.

G9. NIHSS 9. Best language.

- 0 = No aphasia; normal.
- 1 = Mild-to-moderate aphasia; some obvious loss of fluency or facility of comprehension, without significant limitation on ideas expressed or form of expression. Reduction of speech and/or comprehension, however, makes conversation about provided materials difficult or impossible. For example, in conversation about provided materials, examiner can identify picture or naming card content from patient's response.
- 2 = Severe aphasia; all communication is through fragmentary expression; great need for inference, questioning, and guessing by the listener. Range of information that can be exchanged is limited; listener carries burden of communication. Examiner cannot identify materials provided from patient response.
- 3 = Mute, global aphasia; no usable speech or auditory comprehension.
- 3 = Coma/unresponsive. ** An exclusion criterion ** (Choose one answer)

NIHSS 10. Dysarthria.

If patient is thought to be normal, an adequate sample of speech must be obtained by asking patient to read or repeat words from the attached list. If the patient has severe aphasia, the clarity of articulation of spontaneous speech can be rated. Only if the patient is intubated or has other physical barriers to producing speech, the examiner should record the score as untestable (UN), and clearly write an explanation for this choice. Do not tell the patient why he or she is being tested.

G10. NIHSS 10. Dysarthria.

- 0 = Normal.
- 1 = Mild-to-moderate dysarthria; patient slurs at least some words and, at worst, can be understood with some difficulty.
- 2 = Severe dysarthria; patient's speech is so slurred as to be unintelligible in the absence of or out of proportion to any dysphasia, or is mute/anarthric.
- UN = Intubated or other physical barrier. ** An exclusion criterion ** (Choose one answer)

NIHSS 11. Extinction and Inattention (formerly Neglect).

Sufficient information to identify neglect may be obtained during the prior testing. If the patient has a severe visual loss preventing visual double simultaneous stimulation, and the cutaneous stimuli are normal, the score is normal. If the patient has aphasia but does appear to attend to both sides, the score is normal. The presence of visual spatial neglect or anosagnosia may also be taken as evidence of abnormality. Since the abnormality is scored only if present, the item is never untestable.

G11. NIHSS 11. Extinction and Inattention (formerly Neglect).

- 0 = No abnormality.
 1 = Visual, tactile, auditory, spatial, or personal inattention or extinction to bilateral simultaneous stimulation in one of the sensory modalities.
 2 = Profound hemi-inattention or extinction to more than one modality; does not recognize own hand or orients to only one side of space.
 (Choose one answer)

NIHSS & NIHSS-Cog4 scores

G12. Stroke severity as National Institutes of Health stroke scale (NIHSS), NIHSS-Cog4 score: (0-9)

_____ (Calculated)

NIHSS-cog4 = orientation (1b) + executive (1c) + language (9) + inattention (11)

G12a. NIHSS score Stroke severity as National Institutes of Health stroke scale (NIHSS), total 1-11

_____ (Calculated)

△ NIHSS details G1-G11 should be specified.

Section H: Penetration aspiration score (PAS) FEES: Fiberoptic endoscopic evaluation of swallowing**VFS: Videofluoroscopy****PAS: Penetration aspiration score**

H1. Was FEES or VFS performed and PAS measured before baseline/ day 0?

- No FEES or VFS, or no PAS available
 Fiberoptic evaluation of swallowing (FEES)
 Videofluoroscopy (VFS)

H2. Penetration aspiration score (PAS) Specify PAS if performed since treatment on thin fluids (IDDSI level 0) finished using Fiberoptic Evaluation of Swallowing (FEES) or videofluoroscopy (VFS). Please give lowest (best) and highest (worst) PAS if available.

Or give highest (worst) score only if known.

H2a. PAS highest (worst) score

_____ (Integer 1 - 8)

H2b. PAS lowest (best) score

_____ (Integer 1 - 8)

H3. Date PAS score collected

_____ (Date DD-MM-YYYY)

Section I: Glasgow Coma Scale (GCS) - Level of consciousness

I1. Eye Opening Response

4: Spontaneous--open with blinking at baseline
 3: To verbal stimuli, command, speech
 3: To pain only (not applied to face)
 1: No response

I2. Verbal Response

5: Oriented
 4: Confused conversation, but able to answer questions
 3: Inappropriate words
 2: Incomprehensible speech
 1: No response

I3. Motor Response

6 Obeys commands for movement
 5: Purposeful movement to painful stimulus
 4: Withdraws in response to pain
 3: Flexion in response to pain (decorticate posturing)
 2: Extension response in response to pain (decerebrate posturing)
 1: No response

I4a. Glasgow Coma Scale (GCS) total

(Calculated)

△ GCS details I1-I3 should all be specified.

Section J: Assessor information

J1. Please enter your name

(Collected information)

J2a. What is his/her professional role?

- Doctor
 Research coordinator
 Nurse, clinical
 Research nurse
 Physiotherapist
 Occupation therapist
 Speech & Language therapist
 Other
 (Choose one answer)

J2b. If "Other", please specify role

(Professional role)

J3. Does his/her role involve working on stroke wards?

- Yes No
 (Choose one answer)

J4. Please enter your name if you did not collect the information Name of person entering the data, if it differs from the assessor.

(Filling the form)

* Blinded assessors often collect but do not enter the data as it could unblind them.

J5. Please sign the form

(Signature)

Section K: 2nd Assessor information

If there are two members of staff completing the bedside assessment, please complete the second assessor's details below

K1a. Are there two members of staff completing the bedside assessment? Yes No

K1b. 2nd Assessor: Please enter your name?

(Collected information)

K2a. 2nd Assessor: What is your professional role?

- Doctor
 Research coordinator
 Nurse, clinical
 Research nurse
 Physiotherapist
 Occupation therapist
 Speech & Language therapist
 Other
 (Choose one answer)

K2b. If "Other", please specify role

(Professional role)

K3. 2nd Assessor: Does your role involve working on stroke wards? Yes No
(Choose one answer)

K4. 2nd Assessor: Please sign the form

(Signature)

Comments and full explanation for missing data

Are any values missing due to tests not done (or measures not taken), or because data are unknown and every effort has been made to find the data - i.e. 'Not done' / 'Not known'? Yes No

If any values are missing, please provide a full explanation ☐☐ Comments